

## Pandemic Contingency Planning Prioritization Rapid Antigen Test Kits

### Level One (full supply)

- 1) Non-Outbreak sites – staff test 2x week
- 2) Sites in Suspected Outbreak (staff have tested positive, no residents) – staff test daily, people supported test if symptoms appear
- 3) Sites in Outbreak (staff and residents have tested positive) – staff test daily, people supported test if symptoms appear
- 4) Non-Vaccinated Staff test 2x week, at least once at an Agency run Rapid Test Clinic

### Level Two (short supply)

- 1) Non-Outbreak sites – staff test 1x week
- 2) Sites in Suspected Outbreak (staff have tested positive, no residents) *in a program with low numbers of staff* – staff test 2x week, people supported test if symptoms appear
- 3) Sites in Suspected Outbreak (staff have tested positive, no residents) *in a program with a high number of staff* – staff test every other day, people supported test if symptoms appear
- 4) Sites in Outbreak (staff and residents have tested positive) – staff test daily, people supported test if symptoms appear
- 5) Non-Vaccinated Staff test 2x week, at least once at an Agency run Rapid Test Clinic

Any staff with symptoms are not to attend any site.

Staff with symptoms can attend the Agency-run Rapid Test Clinic by providing information ahead of time to the clinic.

**\*\*Health & Safety Committee Members are kept updated on all policies and procedures\*\***